

Preliminary communication

Cyclohexyl(trihalomethyl)mercury compounds: very reactive dihalocarbene precursors

DIETMAR SEYFERTH and CAROL K. HAAS

Department of Chemistry, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139
(U.S.A.)

(Received October 31st, 1972)

SUMMARY

The cyclohexyl(trihalomethyl)mercurials, RHgCCl_3 , RHgCCl_2Br , RHgCClBr_2 and RHgCBr_3 ($\text{R} = \text{cyclohexyl}$), have been prepared and have been found to be excellent dihalocarbene sources. They transfer their CX_2 very rapidly at 80° and within 2–3 days at room temperature.

We have reported recently concerning phenyl(iododichloromethyl)mercury, which is an exceptionally reactive source of dichlorocarbene¹. This reagent transfers CCl_2 to a carbenophile within a few minutes at 80° and within 24 h at room temperature. However, this compound is difficult to prepare and, because of its high reactivity, has poor stability on storage, even as the solid. Phenyl(bromodichloromethyl)mercury transfers CCl_2 fairly rapidly at 80° (within 2 h), but only very slowly at room temperature (within 18 days)². An organomercury CCl_2 transfer reagent with a reactivity intermediate between that of $\text{PhHgCCl}_2\text{I}$ and $\text{PhHgCCl}_2\text{Br}$, one that transfers its CCl_2 very rapidly at 80° and within a few days at 25° , was considered a worthwhile objective for further research.

An indication that such a reagent might be found within the class of the alkyl-(trihalomethyl)mercury compounds was provided by some interesting observations by Shcherbakov³. It was reported that $n\text{-C}_3\text{H}_7\text{HgCCl}_2\text{Br}$ can be isolated as an unpurifiable liquid which decomposes within 2–3 days at room temperature. A three day reaction of this material with cyclohexene at room temperature gave 7,7-dichloronorcarane in 70% yield and $n\text{-C}_3\text{H}_7\text{HgBr}$ (80%). A similar reaction carried out at 40° gave the norcarane in 85% yield within a 5 h period. *N*-Propyl(trichloromethyl)mercury, a liquid, also had enhanced CCl_2 transfer reactivity (compared to PhHgCCl_3), giving 7,7-dichloronorcarane in 82% yield after reaction with cyclohexene at 80° for 8 h. The higher toxicity of alkylmercurials (vs. aryl derivatives) makes work with the relatively volatile *n*-propyl compounds

Base cleavage of trimethylsilyl-1,2-dihydrobenzocyclobutenes, and its significance for the acidities of the 1-, 3-, and 4-positions of 1,2-dihydrobenzocyclobutene ("benzocyclobutene") by C. Eaborn, A. A. Najam and D. R. M. Walton (Brighton) (Rec'd June 7th, 1972)	255
Nuclear magnetic resonance spectra of phenoxysilanes and phenoxygermanes. Evidence for $p\pi-d\pi$ bonding in Si-O and Ge-O bonds by J. R. Chipperfield, D. F. Ewing and G. E. Gould (Hull) (Rec'd June 5th, 1972)	263
Interaction of aryltrimethylstannanes with cyanogen chloride and bromide. A route to aryl cyanides by E. H. Bartlett, C. Eaborn and D. R. M. Walton (Brighton) (Rec'd June 2nd, 1972)	267
Hydrolyses of dimethylchlorotin carboxylates by C. S.-C. Wang and J. M. Shreeve (Moscow, Idaho) (Rec'd June 2nd, 1972)	271
Organometallic compounds. XXXVII. Mass spectra of mixed tetraalkyltins and of trimethyltin halides by M. Gielen and G. Mayence (Brussels) (Rec'd June 7th, 1972)	281
Dipole moments and structure of organophosphine and organoarsine compounds by N. Saraswathi and S. Soundararajan (Bangalore) (Rec'd June 1st, 1972)	289
Alkali-Phosphor-Verbindungen und ihr reaktives Verhalten. LXVII. Bis(diorganylphosphino)-dimethylamino-methane und Diäthylphosphino-dimethylamino-methoxy-methane von K. Issleib und M. Lischewski (Halle/S.) (Eing. den 27. März 1972)	297
Contribution à l'étude des propriétés chimiques des tris(dialcoylamino)stibines par A. Kiennemann, G. Levy et C. Tanielian (Strasbourg) (Reçu le 3 juin 1972)	305
Synthesis and properties of arylidicyclopentadienyltitanium(III) compounds by J. H. Teuben and H. J. de Liefde Meijer (Groningen) (Rec'd June 2nd, 1972)	313
Recherches dans la série des metallocènes. XXIII. Thiophénotropilidènes. Ions et complexes chrome tricarbonyle dérivés. Étude théorique et expérimentale par J. Arriau, J. Deschamps (Pau), T. Marey et J. Tirouflet (Dijon) (Reçu le 3 juin 1972)	323
Intramolecular hydrogen transfer of aminocarbene complexes leading to imines by J. A. Connor and P. D. Rose (Manchester) (Rec'd February 8th, 1972)	329
Organometallic isomerism. The conformation of phosphine-substituted acyldicarbonyl- π -cyclopentadienylmolybdenum complexes in solution by P. J. Craig and J. Edwards (Leicester) (Rec'd May 24th, 1972)	335
Reaction mechanism and properties of the compounds formed in the system: $\text{FeCl}_3 + \text{Mg} + \text{THF} + \text{N}_2$ by B. Jeżowska-Trzebiatowska and P. Sobota (Wrocław) (Rec'd June 2nd, 1972)	339
Infrared spectroscopic studies on metal carbonyl compounds. XIV. Assignments in the C-O stretching region of the authentic infrared spectra of the compounds $\text{MCo}(\text{CO})_9$ ($\text{M} = \text{Mn}$ or Re) and of the monosubstituted derivative $\text{MnCo}(\text{CO})_8\text{P}(\text{C}_6\text{H}_5)_3$ by G. Sbrignadello, G. Bor and L. Maresca (Padova) (Rec'd May 30th, 1972)	345
Infrared spectroscopic studies on metal carbonyl compounds. XV. Assignments of the C-O stretching infrared spectra and force constants of some (arene) $\text{Co}_4(\text{CO})_9$ complexes by G. Bor, G. Sbrignadello and F. Marcati (Padova) (Rec'd May 30th, 1972)	357
Activation and reactivation of catalyst by organic chlorides in the rhodium-catalyzed codimerization of α -olefin and conjugate dienes by A. C. L. Su and J. W. Collette (Wilmington, Del.) (Rec'd April 18th, 1972)	369
The reaction of β -bromostyrene with nickel carbonyl in the presence of alkynes by M. Ryang, Y. Sawa, S. N. Somasundaram, S. Murai and S. Tsutsumi (Osaka) (Rec'd April 14th, 1972)	375
Insertion of sulphur dioxide into platinum-carbon σ -bonds by F. Faraone, L. Silvestro, S. Sergi and R. Pietropaolo (Messina) (Rec'd May 24th, 1972)	379
Autoxidation of arylcopper(I) compounds by A. Camus and N. Marsich (Trieste) (Rec'd July 25th, 1972)	385
Author Index	394
Subject Index	396

















TABLE 1

APPROXIMATE REACTION TIMES FOR CX₂ TRANSFER FROM RHgCX₃

	<i>R</i> = phenyl		<i>R</i> = cyclohexyl	
	room temp.	80°	room temp.	80°
CCl ₃		40 h		8 h
CCl ₂ Br	18 days	2 h	2 days	< 10 min
CClBr ₂	16 days	2 h	3 days	< 10 min
CBr ₃	15 days	2 h	2 days	< 10 min











TABLE 2

REACTIONS OF CYCLOHEXYL (TRIHALOMETHYL) MERCURIALS

CX ₃ in 	Reactant	Reaction conditions ^a	Product (% yield) ^b	 (% yield) ^c
CCl ₃		6.5 h / 80°	 Cl ₂ (92)	
CCl ₃		8 h / 80°	 Cl ₂ (90)	98
CCl ₃	Et ₃ SiH	8 h / 80°	Et ₃ SiCCl ₂ H (88)	97
CCl ₂ Br		6 min / 80°	 Cl ₂ (96)	95
CCl ₂ Br		49 h / 25°	 Cl ₂ (95)	85
CCl ₂ Br		50 h / 25°	 (88)	80
CCl ₂ Br	Et ₃ SiH	50 h / 25°	Et ₃ SiCCl ₂ H (80)	89
CClBr ₂		7 min / 80°	 (85)	90
CClBr ₂		3 days / 25°	 (87)	91

(to be continued)

TABLE 2 (continued)

CX ₃ in 	Reactant	Reaction conditions ^a	Product (% yield) ^b	 (% yield) ^c
CClBr ₂		3 days/25°	 (83)	90
CClBr ₂	Et ₃ SiH	3 days/25°	Et ₃ SiCBrClH (70)	90
CBr ₃		10 min/80°	 (40)	90
CBr ₃		48 h/25°	 (50)	80
CBr ₃		48 h/25°	 (34)	80

^a In general, reactions were carried out with 10 mmol of the mercury reagent and 30 mmol of the reactant in 7–10 ml of dry benzene, with stirring under nitrogen. In room temperature experiments and in the 80° experiments with cyclo-C₆H₁₁HgCCl₃, the benzene solution of the reactants simply was stirred for the stated length of time. In the other 80° experiments, the mercury reagent, dissolved in benzene, was added during 3 min to the preheated substrate and the resulting mixture was heated for the remaining part of the stated reaction time.

^b Determined by GLC.

^c Obtained by filtration of the reaction mixture after it had been cooled to 5°.

undesirable. Furthermore, crystalline, more readily purified alkyl(trihalomethyl)mercurials would be much more practical for synthetic applications. Accordingly, we have searched for alkyl(trihalomethyl)mercury compounds with these desirable properties.

We have found the cyclohexyl(trihalomethyl)mercurials to be reactive dihalo-carbene transfer reagents whose ready application at room temperature makes them particularly attractive. These mercurials were prepared in good yield by the reaction of cyclohexylmercuric chloride, the appropriate haloform and potassium tert-butoxide in THF at -60 to -65°. Their high reactivity requires a rapid work-up of the reaction mixture and storage of the product at 0°. An excess of base must be avoided in these preparations or else pure materials cannot be isolated. These mercurials are subject to autoxidation, hence should be handled under nitrogen. All were obtained as analytically pure, crystalline solids: cyclo-C₆H₁₁HgCCl₃, m.p. 50–52° (lit.⁴ m.p. 52–55°); cyclo-C₆H₁₁HgCCl₂Br, m.p. 44–46°, dec. 85°; cyclo-C₆H₁₁HgCClBr₂, m.p. 60–63°; cyclo-C₆H₁₁HgCBr₃, m.p. 53–56°.


J. Organometal. Chem., 46 (1972)

All of these organomercury reagents except the tribromomethyl compound gave excellent yields of carbene-derived products on decomposition in the presence of a suitable carbenophile in much shorter reaction times than do the analogous phenyl(trihalomethyl)mercurials (Table 1). Each cyclohexyl(trihalomethyl)mercurial was allowed to react with several olefins and with triethylsilane. The results of these experiments are shown in Table 2. Cyclohexyl(trichloromethyl)mercury is shown to be a fairly reactive CCl_2 source at 80° , while cyclohexyl(bromodichloromethyl)mercury reacts very rapidly at 80° but still is quite reactive (compared to $\text{PhHgCCl}_2\text{Br}$) at room temperature. Cyclohexyl(dibromochloromethyl)mercury, a CClBr source, is of comparable reactivity. Unaccountably, cyclohexyl(tribromomethyl)mercury gave good yields of cyclohexylmercuric bromide in comparable reaction times, but the yields of CBr_2 transfer product were not good, at room temperature or at 80° . The formation of a grey color during these cyclo- $\text{C}_6\text{H}_{11}\text{HgCBr}_3$ reactions may indicate an alternate mode of decomposition which does not release dibromocarbene.

An advantage of these reagents is the relatively poor solubility of the cyclohexylmercuric halides in benzene in the concentrations used at room temperature. The cyclohexylmercuric halide yields given in Table 2 represent the material which was filtered from the reaction mixtures after cooling to 5° . At 80° , the cyclohexylmercuric halides were found to be soluble in these reaction mixtures.

These cyclohexyl(trihalomethyl)mercury reagents should prove quite useful in special applications (when the carbenophile is thermally labile or low-boiling or when the product is not thermally stable). It is noteworthy that a relatively reactive organomercury CCl_2 source can now be obtained from the commercially available and inexpensive chloroform.

The acceleration of the rate of CX_2 extrusion from RHgCX_3 compounds when is changed from an aryl group to an alkyl substituent very likely is related to the electron releasing properties of the latter. Transition state I has been suggested for this extrusion process⁵. While aryl substitution on mercury should facilitate nucleophilic attack at



mercury by X, alkyl substitution should assist the heterolysis of the mercury-carbon bond and it appears that the latter is the more important effect.

We are investigating the potential application of the cyclohexyl-for-phenyl variation to some of our other organomercury divalent carbon transfer reagents⁵. Variations in R, the alkyl substituent on mercury, also are being examined.

ACKNOWLEDGMENTS

The authors are grateful to the U.S. Air Force Office of Scientific Research (NC)-

for generous support of this research (Grant AF-AFOSR-72-2204) and to the National Science Foundation for a fellowship to C.K.H.

REFERENCES

- 1 D. Seyferth and C.K. Haas, *J. Organometal. Chem.*, 30 (1971) C38.
- 2 D. Seyferth and H. Shih, *Organometal. Chem. Synth.*, 1 (1972) 41.
- 3 V.I. Shcherbakov, *Zh. Obshch. Khim.*, 41 (1971) 1095.
- 4 G.A. Razuvaev, V.I. Shcherbakov and S.F. Zhil'tsov, *Izv. Akad. Nauk SSSR, Ser. Khim.*, (1968) 2803.
- 5 D. Seyferth, *Accounts Chem. Res.*, 5 (1972) 65.

J. Organometal. Chem., 46 (1972)